

HOSTED BY

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://ees.elsevier.com/ejbas/default.asp>

Full Length Article

Solution of fractional third-order dispersive partial differential equations

Q2

Q1

A.S.V. Ravi Kanth ^{a,*}, K. Aruna ^b^a Department of Mathematics, National Institute of Technology, Kurekshetra 136 119, Haryana, India^b Fluid Dynamics Division, School of Advanced Sciences, VIT University, Vellore, Tamilnadu, India

ARTICLE INFO

Article history:

Received 11 June 2014

Received in revised form

22 November 2014

Accepted 8 February 2015

Available online xxx

Keywords:

Third-order dispersive equations

Fractional differential transform method

Modified fractional differential transform method

ABSTRACT

In this paper, we proposed fractional differential transform method (FDTM) and modified fractional differential transform method (MFDTM) for the solution of fractional third-order dispersive partial differential equations in one- and higher-dimensional spaces. The plotted graphs illustrate the behavior of the solution for different values of fractional order α . The efficiency and accurateness of the proposed methods are examined by means of four numerical experiments.

Copyright 2015, Mansoura University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

In recent past, the glorious developments have been envisaged in the field of fractional calculus and fractional differential equations. Differential equations involving fractional order derivatives are used to model a variety of systems, of which the important applications lie in field of viscoelasticity, electrode-electrolyte polarization, heat conduction, electromagnetic waves, diffusion equation and so on [1,2]. Due to its tremendous scope and applications in several disciplines, a considerable attention has been given to exact and numerical solutions of fractional differential equations. A great deal of

researchers has shown the advantageous use of the fractional calculus in the modeling and control of many dynamical systems [3–10]. Other than modeling aspects of these differential equations, the solution techniques and their reliability are rather more important aspects. It is also equally important to handle critical points which cause sudden divergence convergence and bifurcation of the solutions of the model. In order to achieve the goal of highly accurate and reliable solutions, several methods have been proposed to solve the fractional order differential equations. Some of the recent analytical/numerical methods are Adomian decomposition method (ADM) [11–16], finite difference method [17], Variational iteration method (VIM) [18,19], Operational matrix

* Corresponding author.

E-mail addresses: asvrvikanth@yahoo.com (A.S.V. Ravi Kanth), aruna27k@gmail.com (K. Aruna).

Peer review under responsibility of Mansoura University.

<http://dx.doi.org/10.1016/j.ejbas.2015.02.002>2314-808X/Copyright 2015, Mansoura University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).Please cite this article in press as: Ravi Kanth ASV, Aruna K, Solution of fractional third-order dispersive partial differential equations, Egyptian Journal of Basic and Applied Sciences (2015), <http://dx.doi.org/10.1016/j.ejbas.2015.02.002>

method [20], Homotopy analysis method [21,22], generalized differential transform method [23,24], finite element method [25], fractional differential transform method [26,27] and references therein.

The main aim of this work is to apply FDTM and MFDTM to solve third-order dispersive partial differential equations [28–32]. The DTM obtains an analytical solution in the form of a polynomial. It is different from the traditional high order Taylor's series method, which requires symbolic computation of the necessary derivatives of the data functions. The Taylor series method is computationally taken long time for large orders. With this method, it is possible to obtain highly accurate results or exact solutions for differential equations. The use of DTM in electric circuit analysis was pioneered by Zhou [33]. Since then, DTM was successfully applied for large variety of problems such as partial differential equations [34,35], solitary wave solutions for the KdV and mKdV equations [36], linear and nonlinear Schrodinger equations [37], linear and nonlinear Klein-Gordon equations [38], nonlinear oscillators with fractional nonlinearities [39], fractional linear and nonlinear Schrodinger equation [41], nonlinear fractional Klein-Gordon Equation [42] and references therein. Recently, in [40] presented a novel technique to obtain the differential transform of nonlinearities by the Adomian polynomials. The proposed FDTM and MFDTM do not require linearization, discretization or perturbation unlike the method discussed in the literature. The main drawback of the ADM is to calculate Adomian polynomials for a nonlinear operator where the procedure is very complex. The difficulty in VIM has an inherent inaccuracy in identifying the Lagrange multiplier, correctional functional and stationary conditions for the fractional order. The disadvantage of the Homotopy perturbation method is to solve functional equation in each iteration, which is sometimes complicated and unattainable. Therefore, the proposed FDTM and MFDTM are much easier when compared with ADM, VIM and HPM. The outline of this paper is as follows. In section 2 the basic definitions of fractional calculus are discussed. The basic definitions of two-dimensional FDTM and MFDTM are presented in section three and four. Four clear cut test problems of fractional third-order dispersive partial differential equations are given to elucidate the proposed methods in section 5. At the end, we write the conclusions of the work in section 6.

2. Basic definitions of fractional calculus [2,10]

For convenience of the reader, we present a review of the basic definitions and properties of the fractional calculus theory.

Definition 1. A real function $f(x)$, $x > 0$ in the space C_μ , $\mu \in \mathbb{R}$ if there exists a real number $p > \mu$, such that $f(x) = x^p f_1(x)$, where $f_1(x) \in C[0, \infty)$ and it is said to be in the space iff $f^m \in C_\mu$, $m \in \mathbb{N}$.

Definition 2. The left-sided Riemann-Liouville fractional integral operator of order $\alpha \geq 0$, of a function $f \in C_\mu$, $\mu \geq -1$ is defined as $I^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt$, $\alpha > 0$, $x > 0$ and $J^0 f(x) = f(x)$.

Definition 3. The fractional derivative of $f(x)$ in the Caputo [2] sense is defined as

$$D_*^\alpha f(x) = J^{m-\alpha} D^m f(x) \\ = \frac{1}{\Gamma(m-\alpha)} \int_0^x (x-t)^{m-\alpha-1} f^{(m)}(t) dt \text{ for } m \\ -1 < \alpha < m, m \in \mathbb{N}, x > 0, f \in C^{m-1}.$$

The unknown function $f = f(x, t)$ is assumed to be a casual function of fractional derivatives (i.e., vanishing for $\alpha < 0$) taken in Caputo sense as follows.

Definition 4. For m as the smallest integer that exceeds α , the Caputo time-fractional derivative operator of order $\alpha > 0$ is defined as

$$D_{*t}^\alpha f(x, t) = \frac{\partial^\alpha f(x, t)}{\partial t^\alpha} \\ = \begin{cases} \frac{1}{\Gamma(m-\alpha)} \int_0^t (t-\tau)^{m-\alpha-1} \frac{\partial^m f(x, \tau)}{\partial \tau^m} d\tau, & m-1 < \alpha < m \\ \frac{\partial^m f(x, t)}{\partial t^m}, & \alpha = m \in \mathbb{N} \end{cases}$$

3. Two-dimensional fractional differential transform method

Consider a function of two variables $u(x, t)$ and suppose that it can be represented as a product of two single variable functions i.e., $u(x, t) = f(x)g(t)$. Based on the properties of two-dimensional fractional differential transform, the function $u(x, t)$ can be represented as

$$u(x, t) = \sum_{k=0}^{\infty} \sum_{h=0}^{\infty} U_{\alpha,1}(k, h) (x-x_0)^k (t-t_0)^{h\alpha} \quad (1)$$

where $0 < \alpha$, $U_{\alpha,1}(k, h)$ is called the spectrum of $u(x, t)$. The generalized two-dimensional fractional differential transform of the function $u(x, t)$ is given by

$$U_{\alpha,1} = \frac{1}{\Gamma(k+1)\Gamma(\alpha h+1)} \left[(D_{*x_0}^1)^k (D_{*t_0}^\alpha)^h u(x, t) \right]_{x_0, t_0} \quad (2)$$

where $(D_{*t_0}^\alpha)^h = \underbrace{D_{*t_0}^\alpha D_{*t_0}^\alpha \dots D_{*t_0}^\alpha}_h$. In real applications the function

$u(x, t)$ is represented by a finite series of Eq. (1) can be written as

$$u(x, t) = \sum_{k=0}^l \sum_{h=0}^n U_{\alpha,1}(k, h) x^k t^{h\alpha} + R_{ln}(x, t) \quad (3)$$

and (1) implies that $R_{ln}(x, t) = \sum_{k=l+1}^{\infty} \sum_{h=n+1}^{\infty} U_{\alpha,1}(k, h) x^k t^{h\alpha}$ is negligibly small. Usually, the values of l and n are decided by convergence of the series solution. In case of $\alpha = 1$, the generalized two-dimensional fractional differential transform method (1) reduces to classical two-dimensional differential transform [34–38]. The fundamental mathematical operations performed by two-dimensional fractional differential transform method are listed in Table 1.

Table 1 – The operations for the two-dimensional fractional differential transform method.

Original function	Transformed function
$w(x,t) = u(x,t) \pm v(x,t)$	$W_{\alpha,1}(k,h) = U_{\alpha,1}(k,h) \pm V_{\alpha,1}(k,h)$
$w(x,t) = \mu u(x,t)$	$W_{\alpha,1}(k,h) = \mu U_{\alpha,1}(k,h)$
$w(x,t) = \frac{\partial u(x,t)}{\partial x}$	$W_{\alpha,1}(k,h) = (k+1)U_{\alpha,1}(k+1,h)$
$w(x,t) = D_{t_0^+}^\alpha u(x,t), 0 < \alpha \leq 1$	$W_{\alpha,1}(k,h) = \frac{\Gamma(\alpha(h+1)+1)}{\Gamma(\alpha h+1)} U_{\alpha,1}(k,h+1)$
$w(x,t) = (x-x_0)^m (t-t_0)^{n\alpha}$	$W_{\alpha,1}(k,h) = \delta(k-m, h\alpha-n) = \begin{cases} 1, & k=m, h=n \\ 0, & \text{otherwise} \end{cases}$
$w(x,t) = u^2(x,t)$	$W_{\alpha,1}(k,h) = \sum_{m=0}^k \sum_{n=0}^h U_{\alpha,1}(m,h-n) U_{\alpha,1}(k-m,n)$
$w(x,t) = u^3(x,t)$	$W_{\alpha,1}(k,h) = \sum_{r=0}^k \sum_{q=0}^{k-r} \sum_{s=0}^{h-q} \sum_{p=0}^{h-s-q} U_{\alpha,1}(r,h-s-p) U_{\alpha,1}(q,s) U_{\alpha,1}(k-r-q,p)$

4. Modified fractional differential transform method

However, there are difficulties in FDTM while handling the non-linear functions in two-dimension. Let us consider the differential transform for

$$u^3(x,t) = \sum_{r=0}^k \sum_{q=0}^{k-r} \sum_{s=0}^h \sum_{p=0}^{h-s} U_{\alpha,1}(r,h-s-p) U_{\alpha,1}(q,s) U_{\alpha,1}(k-r-q,p) \tag{4}$$

(4) involves four summations. Thus it is necessary to have a lot of computational work to calculate such differential transform $U_{\alpha,1}(k,h)$ for the large number of (k,h) . As we know that, FDTM is based on the Taylor series for all variables. To avoid these difficulties, MFDTM is considered the Taylor's series of the function $u(x,t)$ with respect to the specific variable. Assume that the specific variable 't' then, we have the Taylor's series expansion of the function $u(x,t)$ at $t=t_0$ as follows.

$$u(x,t) = \sum_{h=0}^{\infty} \frac{1}{\Gamma(\alpha h+1)} \left(\frac{\partial^{\alpha h} u(x,t)}{\partial t^{\alpha h}} \right) (t-t_0)^{\alpha h} \tag{5}$$

Definition 5. The modified fractional differential transform $U_{\alpha,1}(x,h)$ of $u(x,t)$ with respect to the variable t at t_0 is defined by

$$U_{\alpha,1}(x,h) = \frac{1}{\Gamma(\alpha h+1)} \left(\frac{\partial^{\alpha h} u(x,t)}{\partial t^{\alpha h}} \right)_{t=t_0} \tag{6}$$

Definition 6. The modified fractional differential inverse transform $U_{\alpha,1}(x,h)$ of $u(x,t)$ with respect to the variable t at t_0 is defined by

$$u(x,t) = \sum_{h=0}^{\infty} U_{\alpha,1}(x,h) (t-t_0)^{\alpha h} \tag{7}$$

Since the MFDTM results from the Taylor's series of the function with respect to the specific variable it is expected that the corresponding algebraic equation from the given problem is much simpler than the result obtained by the standard FDTM. The fundamental mathematical operations performed by modified fractional differential transform method are listed in Table 2.

5. Applications

In this section, four numerical examples are tested to authenticate the proposed FDTM and MFDTM.

Example 1. We consider the linear fractional dispersive KdV equation

$$u_t^\alpha + 2u_x + u_{xxx} = 0, t > 0 \tag{8}$$

Subject to the initial condition

$$u(x,0) = \sin x \tag{9}$$

FDTM: The transformed version of (8) is

$$\begin{aligned} & \frac{\Gamma(\alpha(h+1)+1)}{\Gamma(\alpha h+1)} U_{\alpha,1}(k,h+1) + 2(k+1)U_{\alpha,1}(k+1,h) \\ & + (k+1)(k+2)(k+3)U_{\alpha,1}(k+3,h) \\ & = 0 \end{aligned} \tag{10}$$

The transformed version of (9) is

$$U_{\alpha,1}(k,0) = \begin{cases} 0, & k = 0, 2, 4, 8, \dots \\ \frac{1}{k!}, & k = 1, 5, 9, \dots \\ -\frac{1}{k!}, & k = 3, 7, 11, \dots \end{cases} \tag{11}$$

Substituting (11) in (10), yields the $U_{\alpha,1}(k,h)$ values,

$$\begin{aligned} U_{\alpha,1}(0,1) &= \frac{-1}{\Gamma(\alpha+1)}, U_{\alpha,1}(1,1) = 0, U_{\alpha,1}(2,1) = \frac{1}{2\Gamma(\alpha+1)}, U_{\alpha,1}(3,1) \\ &= 0, U_{\alpha,1}(4,1) = \frac{-1}{24\Gamma\alpha+1}, \dots \end{aligned}$$

$$\begin{aligned} U_{\alpha,1}(0,2) &= 0, U_{\alpha,1}(1,2) = \frac{-1}{\Gamma(2\alpha+1)}, U_{\alpha,1}(2,2) = 0, U_{\alpha,1}(3,2) \\ &= \frac{1}{6\Gamma(2\alpha+1)}, U_{\alpha,1}(4,2) = 0, \dots \end{aligned}$$

Using $U_{\alpha,1}(k,h)$ values in (1), we obtained the series solution as

$$\begin{aligned} u(x,t) &= \sum_{k=0}^{\infty} \sum_{h=0}^{\infty} U_{\alpha,1}(k,h) x^k t^{\alpha h} \\ &= \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \right) - \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots \right) \frac{t^\alpha}{\Gamma(\alpha+1)} \\ &\quad - \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \right) \frac{t^{2\alpha}}{\Gamma(2\alpha+1)} \\ &\quad + \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots \right) \frac{t^{3\alpha}}{\Gamma(3\alpha+1)} + \dots \end{aligned} \tag{12}$$

When $l,n \leq 5$, then the FDTM solution (3) takes the following form

Table 2 – The operations for the modified fractional differential transform method.

Original function	Transformed function
$w(x,t) = u(x,t) \pm v(x,t)$	$W_{\alpha,1}(x,h) = U_{\alpha,1}(x,h) \pm V_{\alpha,1}(x,h)$
$w(x,t) = \mu u(x,t)$	$W_{\alpha,1}(x,h) = \mu U_{\alpha,1}(x,h)$
$w(x,t) = \frac{\partial u(x,t)}{\partial x}$	$W_{\alpha,1}(x,h) = \frac{\partial U_{\alpha,1}(x,h)}{\partial x}$
$w(x,t) = D_{x_0}^{\alpha} u(x,t), 0 < \alpha \leq 1$	$W_{\alpha,1}(x,h) = \frac{\Gamma(\alpha(h+1)+1)}{\Gamma(\alpha h+1)} U_{\alpha,1}(x,h+1)$
$w(x,t) = (x-x_0)^m (t-t_0)^{n\alpha}$	$W_{\alpha,1}(x,h) = (x-x_0)^m \delta(h\alpha-n)$
$w(x,t) = u^2(x,t)$	$W_{\alpha,1}(x,h) = \sum_{m=0}^h U_{\alpha,1}(x,m) U_{\alpha,1}(x,h-m)$
$w(x,t) = u^3(x,t)$	$W_{\alpha,1}(x,h) = \sum_{m=0}^h \sum_{l=0}^m U_{\alpha,1}(x,h-m) U_{\alpha,1}(x,l) U_{\alpha,1}(x,m-l)$

$$u(x,t) = \left(x - \frac{x^3}{3!} + \frac{x^5}{5!}\right) - \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!}\right) \frac{t^{\alpha}}{\Gamma(\alpha+1)} - \left(x - \frac{x^3}{3!} + \frac{x^5}{5!}\right) \frac{t^{2\alpha}}{\Gamma(2\alpha+1)} + \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!}\right) \frac{t^{3\alpha}}{\Gamma(3\alpha+1)} + \left(x - \frac{x^3}{3!} + \frac{x^5}{5!}\right) \frac{t^{4\alpha}}{\Gamma(4\alpha+1)} - \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!}\right) \frac{t^{5\alpha}}{\Gamma(5\alpha+1)} \quad (13)$$

MFDTM: The transformed version of (8) with respect to 't' is

$$\frac{\Gamma(\alpha(h+1)+1)}{\Gamma(\alpha h+1)} U_{\alpha,1}(x,h+1) + 2 \frac{\partial U_{\alpha,1}(x,h)}{\partial x} + \frac{\partial^3 U_{\alpha,1}(x,h)}{\partial x^3} = 0 \quad (14)$$

The transformed version of (9) is

$$U_{\alpha,1}(x,0) = \sin x \quad (15)$$

The MFDTM recurrence Eq. (14) yields the $U_{\alpha,1}(x,h)$ values

$$U_{\alpha,1}(x,1) = \frac{-\cos x}{\Gamma(\alpha+1)}, U_{\alpha,1}(x,2) = \frac{-\sin x}{\Gamma(2\alpha+1)}, U_{\alpha,1}(x,3) = \frac{\cos x}{\Gamma(3\alpha+1)}, \dots$$

Substituting $U_{\alpha,1}(x,h)$'s into Eq. (7), we obtained solution in the following form

$$u(x,t) = \sin x - \frac{\cos x}{\Gamma(\alpha+1)} t^{\alpha} - \frac{\sin x}{\Gamma(2\alpha+1)} t^{2\alpha} + \dots \quad (16)$$

When $\alpha \rightarrow 1$, the approximate solution Eqs. (12) and (16) takes the following form

$$u(x,t) = \sin x \cos t - \cos x \sin t = \sin(x-t)$$

which is exactly same as the solution obtained in [32]. In Figs. 1–2 we have shown the solution $u(x,t)$ obtained by five term FDTM, MFDTM and exact solution for the two different values of fractional order α . It can be seen that the MFDTM results are far better approximations than the FDTM results.

Example 2. Next, consider the linear fractional dispersive KdV equation in two-dimensional space

$$u_t^{\alpha} + 2u_{xxx} + u_{yyy} = 0, t > 0 \quad (17)$$

Subject to the initial condition

$$u(x,y,0) = \cos(x+y) \quad (18)$$

FDTM: The transformed version of (17) is

$$\frac{\Gamma(\alpha(h+1)+1)}{\Gamma(\alpha h+1)} U_{\alpha,1}(k,l,h+1) + (k+1)(k+2)(k+3) U_{\alpha,1}(k+3,l,h) + (l+1)(l+2)(l+3) U_{\alpha,1}(k,l+3,h) = 0 \quad (19)$$

The transformed version of (18) is

$$U_{\alpha,1}(k,l,0) = \frac{\cos\left(\frac{k\pi}{2}\right) \cos\left(\frac{l\pi}{2}\right) \sin\left(\frac{k\pi}{2}\right) \sin\left(\frac{l\pi}{2}\right)}{k! l!}, k, l = 0, 1, 2, \dots \quad (20)$$

Substituting Eq. (20) in Eq. (19), yields the $U_{\alpha,1}(k,l,h)$ values

$$U_{\alpha,1}(0,0,1) = 0, U_{\alpha,1}(1,0,1) = \frac{-2}{\Gamma(1+\alpha)}, U_{\alpha,1}(2,0,1) = 0, U_{\alpha,1}(3,0,1) = \frac{1}{3\Gamma(1+\alpha)}, \dots$$

$$U_{\alpha,1}(0,1,1) = \frac{-2}{\Gamma(1+\alpha)}, U_{\alpha,1}(1,1,1) = 0, U_{\alpha,1}(2,1,1) = 0, U_{\alpha,1}(3,1,1) = 0, \dots$$

$$U_{\alpha,1}(0,2,1) = 0, U_{\alpha,1}(1,2,1) = \frac{1}{\Gamma(1+\alpha)}, U_{\alpha,1}(2,2,1) = 0, U_{\alpha,1}(3,2,1) = \frac{-1}{6\Gamma(1+\alpha)}, \dots$$

Using $U_{\alpha,1}(k,l,h)$ values in Eq. (1), we obtained the series solution as

$$u(x,y,t) = 1 - \frac{x^2}{2} - xy - \frac{x^3 y}{6} - \frac{y^2}{2} + \frac{x^2 y^2}{4} + \frac{xy^3}{6} - \frac{x^3 y^3}{36} + \dots + \left(-2x + \frac{x^3}{3} - 2y + x^2 y + xy^2 - \frac{x^3 y^2}{6} + \frac{y^3}{3} - \frac{x^2 y^3}{6} + \dots\right) \frac{t^{\alpha}}{\Gamma(1+\alpha)} + \dots \quad (21)$$

MFDTM: The transformed version of Eq. (17) is w. r. t 't' is

$$\frac{\Gamma(\alpha(h+1)+1)}{\Gamma(\alpha h+1)} U_{\alpha,1}(x,y,h+1) + \frac{\partial^3 U_{\alpha,1}(x,y,h)}{\partial x^3} + \frac{\partial^3 U_{\alpha,1}(x,y,h)}{\partial y^3} = 0 \quad (22)$$

The transformed version of (18) is

$$U_{\alpha,1}(x,y,0) = \cos(x+y) \quad (23)$$

The MFDTM recurrence Eq. (22) yields the $U_{\alpha,1}(x,y,h)$ values




Fig. 1 – $u(x,t)$ obtained by using five term (a) FDTM, (b) MFDTM and (c) Exact solution when $\alpha = 0.95$.

$$U_{\alpha,1}(x, y, 1) = \frac{-2 \sin(x+y)}{\Gamma(\alpha+1)}, U_{\alpha,1}(x, y, 2) = \frac{-4 \cos(x+y)}{\Gamma(2\alpha+1)}, U_{\alpha,1}(x, y, 3) = \frac{8 \sin(x+y)}{\Gamma(3\alpha+1)}, \dots$$

Using the inverse MFDTM, we obtained the solution in the following form,

$$u(x, y, t) = \cos(x+y) - \frac{2 \sin(x+y)}{\Gamma(\alpha+1)} t^\alpha - \frac{4 \cos(x+y)}{\Gamma(2\alpha+1)} t^{2\alpha} + \dots \quad (24)$$

When $\alpha \rightarrow 1$, the approximate solution (21) and (24) takes the following form

$$u(x, t) = \sin(x+y+2t)$$

which is exactly same as the solution obtained in [32].

Example 3. Consider the non-homogeneous fractional third-order dispersive partial differential equation

$$u_t^\alpha + u_{xxx} = -\sin \pi x \sin t - \pi^3 \cos \pi x \cos t, \quad 0 < x < 1, t > 0 \quad (25)$$

Subject to the initial condition

$$u(x, 0) = \sin \pi x \quad (26)$$

and time-dependent boundary conditions

$$u(0, t) = 0, u_x(0, t) = \pi \cos t, u_{xx}(0, t) = 0 \quad (27)$$

FDTM: The transformed version of (25) is

$$\begin{aligned} & \frac{\Gamma(\alpha(h+1)+1)}{\Gamma(\alpha h+1)} U_{\alpha,1}(k, h+1) + (k+1)(k+2)(k+3)U_{\alpha,1}(k+3, h) \\ & = -(\pi)^k \frac{\sin\left(\frac{k\pi}{2}\right)}{k!} \frac{\sin\left(\frac{h\pi}{2}\right)}{\Gamma(\alpha h+1)} - \pi^3 (\pi)^k \frac{\cos\left(\frac{k\pi}{2}\right)}{k!} \frac{\cos\left(\frac{h\pi}{2}\right)}{\Gamma(\alpha h+1)} \end{aligned} \quad (28)$$

The transformed version of (26) is

$$U_{\alpha,1}(k, 0) = \begin{cases} 0, & k = 0, 2, 4, 6, \dots \\ \frac{(\pi)^k}{k!}, & k = 1, 5, 9, \dots \\ -\frac{(\pi)^k}{k!}, & k = 3, 7, 11, \dots \end{cases} \quad (29)$$

MFDTM: The transformed version of (25) is w. r. t 't' is




Fig. 2 – $u(x,t)$ obtained by using five term (a) FDTM, (b) MFDTM and (c) Exact solution when $\alpha = 1.5$.

$$\frac{\Gamma(\alpha(h+1)+1)}{\Gamma(\alpha h+1)} U_{\alpha,1}(x, h+1) + \frac{\partial^3 U_{\alpha,1}(x, h)}{\partial x^3} = \sin \pi x \frac{\sin\left(\frac{\pi h}{2}\right)}{\Gamma(\alpha h+1)} - \pi^3 \cos \pi x \frac{\cos\left(\frac{\pi h}{2}\right)}{\Gamma(\alpha h+1)} \quad (30)$$

The transformed version of (26) is

$$U_{\alpha,1}(x, 0) = \sin \pi x \quad (31)$$

Following the same procedure in Example 1 and 2, we obtained the FDTM and MFDTM series solution

$$u(x, t) = \left(\pi x - \frac{(\pi x)^3}{3!} + \frac{(\pi x)^5}{5!} - \dots \right) \left(1 - \frac{t^{2\alpha}}{\Gamma(2\alpha+1)} + \frac{t^{4\alpha}}{\Gamma(4\alpha+1)} - \dots \right) \quad (32)$$

$$u(x, t) = \sin \pi x \left(1 - \frac{t^{2\alpha}}{\Gamma(2\alpha+1)} + \frac{t^{4\alpha}}{\Gamma(4\alpha+1)} - \dots \right) \quad (33)$$

respectively. When $\alpha \rightarrow 1$, the approximate solution (32) and (33) takes the following form

$$u(x, t) = \sin \pi x \cos t \quad (34)$$

which is exactly same as the solution obtained in [32]. Five term solution of $u(x,t)$ using FDTM, MFDTM corresponding to the values of $\alpha=0.5, 1.95$ and exact solution are plotted in Figs. 3–4 respectively.

Example 4. Finally, consider the non-homogeneous fractional third-order dispersive partial differential equation in three dimensional space

$$u_t^\alpha + u_{xxx} + \frac{1}{8} u_{yyy} + \frac{1}{27} u_{zzz} = -3 \cos(x+2y+3z) \sin t + \sin(x+2y+3z) \cos t, t > 0 \quad (35)$$

Subject to the initial condition

$$u(x, y, z, 0) = 0 \quad (36)$$

FDTM: The transformed version of (35) is

$$\begin{aligned}
 & \frac{\Gamma(\alpha(h+1)+1)}{\Gamma(\alpha h+1)} U_{\alpha,1}(k,l,m,h+1) + (k+1)(k+2)(k+3)U_{\alpha,1}(k+3,l,m,h) \\
 & + \frac{1}{8}(l+1)(l+2)(l+3)U_{\alpha,1}(k,l+3,m,h) + \frac{1}{27}(m+1)(m+2)(m+3)U_{\alpha,1}(k,l,m+3,h) \\
 & = \left(-3 \frac{\cos\left(\frac{k\pi}{2}\right)}{k!} \frac{(3)^m \cos\left(\frac{m\pi}{2}\right)}{m!} + 3 \frac{\sin\left(\frac{k\pi}{2}\right)}{k!} \frac{(3)^m \sin\left(\frac{m\pi}{2}\right)}{m!} \right) \frac{(2)^l \cos\left(\frac{l\pi}{2}\right) \sin\left(\frac{h\pi}{2}\right)}{l! \Gamma(\alpha h+1)} \\
 & + \left(3 \frac{\sin\left(\frac{k\pi}{2}\right)}{k!} \frac{(3)^m \cos\left(\frac{m\pi}{2}\right)}{m!} + 3 \frac{\cos\left(\frac{k\pi}{2}\right)}{k!} \frac{(3)^m \sin\left(\frac{m\pi}{2}\right)}{m!} \right) \frac{(2)^l \sin\left(\frac{l\pi}{2}\right) \sin\left(\frac{h\pi}{2}\right)}{l! \Gamma(\alpha h+1)} \\
 & + \left(\frac{\sin\left(\frac{k\pi}{2}\right)}{k!} \frac{(3)^m \cos\left(\frac{m\pi}{2}\right)}{m!} - \frac{\cos\left(\frac{k\pi}{2}\right)}{k!} \frac{(3)^m \sin\left(\frac{m\pi}{2}\right)}{m!} \right) \frac{(2)^l \cos\left(\frac{l\pi}{2}\right) \cos\left(\frac{h\pi}{2}\right)}{l! \Gamma(\alpha h+1)} \\
 & - \left(\frac{\cos\left(\frac{k\pi}{2}\right)}{k!} \frac{(3)^m \cos\left(\frac{m\pi}{2}\right)}{m!} - \frac{\sin\left(\frac{k\pi}{2}\right)}{k!} \frac{(3)^m \sin\left(\frac{m\pi}{2}\right)}{m!} \right) \frac{(2)^l \sin\left(\frac{l\pi}{2}\right) \cos\left(\frac{h\pi}{2}\right)}{l! \Gamma(\alpha h+1)}
 \end{aligned} \tag{37}$$




Fig. 3 – $u(x,t)$ obtained by using five term (a) FDTM, (b) MFDTM and (c) Exact solution when $\alpha = 0.5$.




Fig. 4 – $u(x,t)$ obtained by using five term (a) FDTM, (b) MFDTM and (c) Exact solution when $\alpha = 1.95$.

The transformed version of (36) is

$$U_{\alpha,1}(k, l, m, 0) = 0, k, l, m = 0, 1, 2, \dots \tag{38}$$

MFDTM: The transformed version of (35) is w. r. t 't' is

$$\begin{aligned} & \frac{\Gamma(\alpha(h+1)+1)}{\Gamma(\alpha h+1)} U_{\alpha,1}(x, y, z, h+1) + \frac{\partial^3 U_{\alpha,1}(x, y, z, h)}{\partial x^3} \\ & + \frac{1}{8} \frac{\partial^3 U_{\alpha,1}(x, y, z, h)}{\partial y^3} + \frac{1}{27} \frac{\partial^3 U_{\alpha,1}(x, y, z, h)}{\partial z^3} \\ & = -3 \cos(x+2y+3z) \frac{\sin\left(\frac{\pi h}{2}\right)}{\Gamma(\alpha h+1)} + \sin(x+2y+3z) \frac{\cos\left(\frac{\pi h}{2}\right)}{\Gamma(\alpha h+1)} \end{aligned} \tag{39}$$

The transformed version of (36) is

$$U_{\alpha,1}(x, y, z, 0) = 0 \tag{40}$$

Following the same procedure in Example 1 and 2, we obtained the FDTM and MFDTM series solution

$$\begin{aligned} u(x, t) = & \left(x + 2y + 3z - \frac{x^3}{6} - \frac{4}{3}y^3 - \frac{9}{2}z^3 - x^2y - 9z^2y - \frac{3}{2}zx^2 \right. \\ & \left. - 2y^2x - 6zy^2 + \dots \right) \frac{t^\alpha}{\Gamma(\alpha+1)} - \dots \end{aligned} \tag{41}$$

$$u(x, y, z, t) = \sin(x+2y+3z) \left(\frac{t^\alpha}{\Gamma(\alpha+1)} - \frac{t^{3\alpha}}{\Gamma(3\alpha+1)} + \dots \right) \tag{42}$$

respectively. When $\alpha \rightarrow 1$, the approximate solution (32) and (33) takes the following form

$$u(x, y, z, t) = \sin(x+2y+3z) \sin t \tag{43}$$

which is exactly same as the solution obtained in [32].

6. Conclusions

In this paper, we implemented the two-dimensional FDTM and MFDTM for solving fractional third-order dispersive partial differential equation. DTM is an attractive tool for solving linear and nonlinear partial differential equations and it does

not require linearization, discretization or perturbation. But it also faces some difficulties while constructing recursive equation for the function of three or more variables and it requires an expensive computational cost to solve the algebraic recursive equation. The proposed MFDTM for the specific variable can obtain the simple recursive equation. Thus it is concluded that MFDTM enhances the effectiveness of the computational work when compared with the FDTM. The proposed methods are simpler in its principles and effective in solving linear and nonlinear differential equations of fractional order and promising tool for solving wider class of nonlinear fractional models in mathematical physics with high accuracy.

Acknowledgments

The authors would like to thank the referees for their comments and suggestions that have improved the paper.

REFERENCES

- [1] Hilfer R. Applications of fractional calculus in physics. Singapore: Word Scientific Company; 2000.
- [2] Caputo M. Linear models of dissipation whose Q is almost frequency independent. Part II Geophys J RAstron Soc 1967;13:529–39.
- [3] Kilbas Anatoly A, Srivastava HM, Trujillo Juan J. Theory and applications of fractional differential equations. North-Holland: Jan Van Mill; 2006.
- [4] Petras Ivo. Fractional-order nonlinear systems: modeling, analysis and simulation. Beijing: Higher Education Press; 2011.
- [5] Laroche E, Knittel D. An improved linear fractional model for robustness analysis of a winding system. Control Eng Pract 2005;13:659–66.
- [6] Calderon A, Vinagre B, Feliu V. Fractional order control strategies for power electronic buck converters. Signal Process 2006;86:2803–19.
- [7] Sabatier J, Aoun M, Oustaloup A, Grgoire G, Ragot F, Roy P. Fractional system identification for lead acid battery state of charge estimation. Signal Process 2006;86:2645–57.
- [8] Vinagre B, Monje C, Calderon A, Suarez J. Fractional PID controllers for industry application. A brief introduction. J Vib Control 2007;13:1419–30.
- [9] Monje C, Vinagre B, Feliu V, Chen Y. Tuning and autotuning of fractional order controllers for industry applications. Control Eng Pract 2008;16:798–812.
- [10] Podlubny I. Fractional differential equations. San Diego: Academic Press; 1999.
- [11] Ray SS, Bera RK. Analytical solution of Bagley–Torvik equation by Adomian decomposition method. Appl Math Comput 2005;168(1):398–410.
- [12] Momani S, Odibat Z. Numerical approach to differential equations of fractional orders. J Comput Appl Math 2007;207(1):96–110.
- [13] Duan Jun-sheng, Jianye An, Mingyu Xu. Solution of system of fractional differential equations by Adomian decomposition method. Appl Mathematics-A J Chin Univ 2007;22(1):7–12.
- [14] Odibat Z, Momani S. Numerical methods for nonlinear partial differential equations of fractional order. Appl Math Model 2008;32:28–9.
- [15] Li Changpin, Wang Yihong. Numerical algorithm based on Adomian decomposition for fractional differential equations. Comput Math Appl 2009;57(10):1672–81.
- [16] Duan Jun-sheng, Chaolu Temuer, Rach Randolph, Lu Lei. The Adomian decomposition method with convergence acceleration techniques for nonlinear fractional differential equations. Comput Math Appl 2013;66(5):728–36.
- [17] Meerschaert M, Tadjeran C. Finite difference approximations for two sided space fractional partial differential equations. Appl Numer Math 2006;56:80–90.
- [18] Mustafa Inc. The approximate and exact solutions of the space- and time- fractional Burger's equations with initial conditions by variational iteration method. J Math Anal Appl 2008;345(1):476–84.
- [19] Das S. Analytical solution of a fractional diffusion equation by variational iteration method. Comput Math Appl 2009;57:483–7.
- [20] Saadatmandi A, Dehghan M. A new operational matrix for solving fractional order differential equations. Comput Math Appl 2010;59:1326–36.
- [21] Pandey RK, Singh OP, Baranwal VK. An analytic algorithm for the space–time fractional advection–dispersion equation. Comput Phys Commun 2011;182:1134–44.
- [22] Khan Najeeb Alam, Khan Nasir-uddin, Ara Asmat, Jamil Muhammad. Approximate analytical solutions of fractional reaction-diffusion equations. J King Saud Univ – Sci 2012;24(2):111–8.
- [23] Odibat Z, Momani S, Erturk VS. Generalized differential transform method: application to differential equations of fractional order. Appl Math Comput 2008;197:467–77.
- [24] Liu J, Hou G. Numerical solutions of the space- and timefractional coupled Burgers equation by generalized differential transform method. Appl Math Comput 2011;217(16):7001–8.
- [25] Jiang Y, Ma J. Higher order finite element methods for time fractional partial differential equations. J Comput Appl Math 2011;235(11):3285–90.
- [26] Arikoglu A, Ozkol I. Solution of a fractional differential equations by using differential transform method. Chaos Solit Fractals 2007;34:1473–81.
- [27] Erturk VS, Momani S. Solving systems of fractional differential equations using differential transform method. J Comput Appl Math 2008;215:142–51.
- [28] Djidjeli K, Twizell EH. Global extrapolations of numerical methods for solving a third-order dispersive partial differential equations. Int J Comput Math 1991;41:81–9.
- [29] Twizell EH. Computational methods for partial differential equations. Ellis Horwood, New York: Chichester and John Wiley and Sons; 1984.
- [30] Lewson JD, Morris JLI. The extrapolation of first-order methods for parabolic partial differential equations I. SIAM J Numer Anal 1978;15:1212–24.
- [31] Mengzhaoh Q. Difference scheme for the dispersive equation. Computing 1983;31:261–7.
- [32] Wazwaz AM. An analytic study on the third-order dispersive partial differential equation. Appl Math Comput 2003;142:511–20.
- [33] Zhou JK. Differential transformation and its applications for electrical circuits. Wuhan: Huazhong University Press; 1986.
- [34] Chen CK, Ho SH. Solving partial differential equations by two-dimensional differential transform method. Appl Math Comput 1999;106:171–9.
- [35] Jang MJ, Chen CL, Liu YC. Two-dimensional differential transform for partial differential equations. Appl Math Comput 2001;121:261–70.
- [36] Figen Kangalgil O, Ayaz F. Solitary wave solutions for the KdV and mKdV equations by differential transform method. Chaos Solit Fractals 2009;41(1):464–72.

- 1 [37] Ravi Kanth ASV, Aruna K. Two-dimensional differential
2 transform method for solving linear and non-linear
3 Schrodinger equations. *Chaos Solit Fractals* 2009;41:2277–81.
- 4 [38] Ravi Kanth ASV, Aruna K. Differential transform method for
5 solving the linear and nonlinear Klein–Gordon equation.
6 *Comput Phys Commun* 2009;180:708–11.
- 7 [39] Ebaid AE. A reliable after treatment for improving the
8 differential transformation method and its application to
9 nonlinear oscillators with fractional nonlinearities. *Commun
10 Nonlinear Sci Numer Simul* 2011;16:528–36.
- 11 [40] Fatoorehchi H, Abolghasemi H. Improving the differential
transform method: a novel technique to obtain the
differential transforms of nonlinearities by the Adomian
polynomials. *Appl Math Model* 2013;37:6008–17.
- [41] Aruna K, Ravi Kanth ASV. Approximate solutions of non-
linear fractional Schrodinger equation via differential
transform method and modified differential transform
method. *Natl Acad Sci Lett* 2013;36(2):201–13.
- [42] Aruna K, Ravi Kanth ASV. Two-dimensional differential
transform method and modified differential transform
method for solving nonlinear fractional Klein-Gordon
equation. *Natl Acad Sci Lett* 2014;37(2):163–71.

UNCORRECTED PROOF